



ATLAS VETERINAIRE
Laboratoire pharmaceutique vétérinaire

ERYTHROFORT

ORAL POWDER

COMPOSITION

Erythromycin (sf. Thiocyanate) 20.00 g
Excipient qs 100 g

PHARMACEUTICAL PROPERTIES

Erythromycin is an antibiotic from the macrolide family with great activity, especially on gram-positive germs and mycoplasmas. Erythromycin is obtained by culturing a strain of streptomyces erythreus. It is a bacteriostatic antibiotic. Its mode of action results in the inhibition of bacterial protein synthesis. Erythromycin is well absorbed from the digestive tract in monogastrics and is widely distributed in the lung, kidney, liver and skin. Erythromycin is eliminated primarily via the bile after hepatic metabolism.

TARGET SPECIES

Poultry.

INDICATIONS

Infections with germs sensitive to Erythromycin. It is mainly concerned with the treatment and prevention of avian mycoplasmosis (primary and CKD), infectious coryza, infectious sinusitis and synovitis in poultry.

ADMINISTRATION AND DOSAGE

Oral use after dilution in drinking water the dose of: 1.0 g / liter of drinking water, or a sachet of 1 kg / ton of water. The duration of treatment is 5 days. It is recommended to pre-dilute the contents of the sachet in 10 to 20 liters of water. Avoid any other source of water for the duration of treatment.

CONTRAINDICATIONS AND INTERACTIONS

Hypersensitivity to Erythromycin. Hepatic dysfunction. Do not administer to rabbits. Erythromycin can be safely combined with several other antibacterials such as: tetracyclines, aminoglycosides and sulfonamides.



ATLAS VÉTÉRINAIRE
Laboratoire pharmaceutique vétérinaire

SIDE EFFECTS

If the doses and contraindications are respected, no side effects are observed in the target species.

PRECAUTIONS FOR USE

Respect the dosages. Keep the product out of the reach of children.

Keep the product at a temperature not exceeding 25 ° C and protect from light.

WITHDRAWAL PERIOD

Meat and offal: 3 days Eggs: 2 days

STORAGE CONDITIONS

Keep the product in the original packaging, protect from light and at a temperature ≤ 25 ° C.

PHARMACEUTICAL FORM AND PRESENTATION

Sachets of 500 g and 1 Kg