



ATLAS VETERINAIRE
Laboratoire pharmaceutique vétérinaire

ZURICOX

ORAL SOLUTION

COMPOSITION

Toltrazuril2,50 g
Excipient q.s.p.....100 ml

PHARMACEUTICAL PROPERTIES

Toltrazuril, active ingredient in the specialty ZURICOX, is an anticoccidial from the triazinone group active on all intracellular stages of coccidial development. Toltrazuril acts on schizonts (schizogony) and micro- and macrogametes (gametogony). Toltrazuril interferes with the division of the coccidial nucleus and the activity of mitochondria, and induces severe vacuolation of the protoplasmic endoplasmic reticulum. Toltrazuril does not interfere with the development of anticoccidial immunity, and its effectiveness is independent of the severity of the infection. Toltrazuril is also active against *Toxoplasma*, *Theileria* and *sarcocystis*. In the body, toltrazuril is eliminated mainly in the faeces in unchanged or oxidized form (sulfoxide and sulfone).

TARGET SPECIES

Poultry.

INDICATIONS

Prevention and treatment of avian coccidiosis due to *Eimeria acervulina*, *E. necatrix*, *E. brunetti*, *E. maxima* and *E. mitis* in chicken and to *E. adenoides* and *E. meleagridis* in turkeys.

ADMINISTRATION AND DOSAGE

Oral use in drinking water at the following dosage: Basic dosage: 7 mg of active ingredient / kg bodyweight / day, i.e. 28 ml of ZURICOX solution per 100 kg per day for 2 consecutive days. Prevention: before high-risk periods (stress, vaccination, transport) in strategic administration of 2 consecutive days. Treatment: administration for 2 consecutive days in the presence or absence of anticoccidians in the food.



ATLAS VETERINAIRE
Laboratoire pharmaceutique vétérinaire

CONTRAINDICATIONS AND INTERACTIONS

Do not use in laying hens whose eggs or their derivatives are intended for human consumption.

SIDE EFFECTS

If the general conditions of use are respected, no undesirable effects are observed after treatment with Toltrazuril.

PRECAUTIONS FOR USE

- Respect the indicated dosage.
- Shake well before use.
- Wash hands thoroughly after using the product.
- Avoid contact with the skin; In case of accidental ingestion, consult a doctor.
- Keep the product out of reach of children.

WITHDRAWAL PERIOD

- Meat and offal: 12 days
- Milk: not applicable
- Eggs: Do not use in laying hens whose eggs or their derivatives are intended for human consumption

STORAGE CONDITIONS

Keep the product in the original packaging, protect from light and at a temperature ≤ 25 ° C.

PHARMACEUTICAL FORM AND PRESENTATION

500 ml and 1-liter bottles