



# EQUIPAZ

## ORAL SUSPENSION

### COMPOSITION

Ivermectin .....	0.50 g
Praziquantel .....	2.50 g
Excipient qs .....	100 ml

### PHARMACEUTICAL PROPERTIES

Ivermectin, an antiparasitic of the family of avermectins endowed with a great activity as well on the internal parasites as external of the domestic animals. Ivermectin acts on nematodes and arthropods via gamma amino-butyric acid (GABA). It causes its release from the presynaptic membrane and potentiates its binding to receptors located on the postsynaptic membrane. At the recommended doses, ivermectin is very well tolerated in both adults and young people and can be used at all stages of gestation. However, ivermectin is devoid of cestocidal activity. This often forces it to be combined with anthelmintics with tenicidal activity.

Praziquantel is an anthelmintic from the pyrazinoisoquinoline family which is very active on tapeworms. Its action is exerted on both adult and immature forms. On the pharmacological level, praziquantel exerts its action in various ways including in particular the induction of a paralyzing muscular contraction and the destruction of the integument of the cestode. It is rapidly absorbed from the digestive tract in all species. His metabolism is complete and rapid. The two products, ivermectin and praziquantel, then present a perfect complementarity of their action spectra.



### TARGET SPECIES

Equines

### INDICATIONS

Treatment of gastrointestinal and pulmonary nematodoses and Cestodoses in equines.



**ATLAS VETERINAIRE**  
Laboratoire pharmaceutique vétérinaire

### **ADMINISTRATION AND DOSAGE**

EQUIPAZ is recommended orally in a single dose of 4.00 ml suspension / 100 kg bodyweight, corresponding to 200 µg ivermectin and 1 mg praziquantel per kg bodyweight.

### **CONTRAINDICATIONS AND INTERACTIONS**

Do not use in foals under 5 months of age.

Do not use in slaughter horses

### **SIDE EFFECTS**

If the general conditions of use are respected, no side effects are observed.

### **PRECAUTIONS FOR USE**

- Do not smoke or eat while using the product.
- Wash hands thoroughly after using the product.
- Keep the product out of the reach of children.
- Destroy the bottle after use.

### **WITHDRAWAL PERIOD**

Do not use in slaughter horses

### **STORAGE CONDITIONS**

Keep the product in the original packaging, protect from light and at a temperature  $\leq 25^{\circ}$  C.

### **PRESENTATION**

S Bottles of 20 ml and 50 ml.